


MEGESTROL ACETATE - megestrol acetate tablet

Roxane Laboratories, Inc

RX ONLY

DESCRIPTION

Megestrol acetate, is a synthetic, antineoplastic and progestational drug. Megestrol acetate is a white, crystalline solid chemically designated as pregna-4,6-diene-3,20-dione, 17-(acetyloxy)-6-methyl-. Solubility at 37°C in water is 2 mcg per mL, solubility in plasma is 24 mcg per mL. Its molecular weight is 384.51. The empirical formula is C₂₄ H₃₂ O₄ and the structural formula is represented as follows:



Megestrol acetate tablets are supplied for oral administration containing 20 mg or 40 mg megestrol acetate and the following inactive ingredients: acacia, colloidal silicon dioxide, dibasic calcium phosphate, lactose monohydrate, magnesium stearate, and starch (corn).

CLINICAL PHARMACOLOGY

While the precise mechanism by which megestrol produces its antineoplastic effects against endometrial carcinoma is unknown at the present time, inhibition of pituitary gonadotrophin production and resultant decrease in estrogen secretion may be factors. There is evidence to suggest a local effect as a result of the marked changes brought about by the direct instillation of progestational agents into the endometrial cavity. The antineoplastic action of megestrol acetate on carcinoma of the breast is effected by modifying the action of other steroid hormones and by exerting a direct cytotoxic effect on tumor cells. In metastatic cancer, hormone receptors may be present in some tissues but not others. The receptor mechanism is a cyclic process whereby estrogen produced by the ovaries enters the target cell, forms a complex with cytoplasmic receptor and is transported into the cell nucleus. There it induces gene transcription and leads to the alteration of normal cell functions. Pharmacologic doses of megestrol acetate not only decrease the number of hormone-dependent human breast cancer cells but also are capable of modifying and abolishing the stimulatory effects of estrogen on these cells. It has been suggested that progestins may inhibit in one of two ways: by interfering with either the stability, availability, or turnover of the estrogen receptor complex in its interaction with genes or in conjunction with the progestin receptor complex, by interacting directly with the genome to turn off specific estrogen-responsive genes.

There are several analytical methods used to estimate megestrol acetate plasma levels, including mass fragmentography, gas chromatography (GC), high pressure liquid chromatography (HPLC) and radioimmunoassay. The plasma levels by HPLC assay or radioimmunoassay methods are about one-sixth those obtained by the GC method. The plasma levels are dependent not only on the method used, but also on intestinal and hepatic inactivation of the drug, which may be affected by factors such as intestinal tract motility, intestinal bacteria, antibiotics administered, body weight, diet, and liver function.

Metabolites account for only 5% to 8% of the administered dose and are considered negligible. The major route of drug elimination in humans is the urine. When radiolabeled megestrol acetate was administered to humans in doses of 4 to 90 mg, the urinary excretion within 10 days ranged from 56.5% to 78.4% (mean 66.4%) and fecal excretion ranged from 7.7% to 30.3% (mean 19.8%). The total recovered radioactivity varied between 83.1% and 94.7% (mean 86.2%). Respiratory excretion as labeled carbon dioxide and fat storage may have accounted for at least part of the radioactivity not found in the urine and feces.

In normal male volunteers (n=23) who received 160 mg of megestrol acetate given as a 40 mg q.i.d. regimen, the oral absorption of megestrol acetate appeared to be variable. Plasma levels were assayed by a high pressure liquid chromatographic (HPLC) procedure. Peak drug levels for the first 40 mg dose ranged from 10 to 56 ng/mL (mean 27.6 ng/mL) and the times to peak concentrations ranged from 1.0 to 3.0 hours (mean 2.2 hours). Plasma elimination half-life ranged from 13.0 to 104.9 hours (mean 34.2 hours). The steady state plasma concentrations for a 40 mg q.i.d. regimen have not been established.

INDICATIONS AND USAGE

Megestrol acetate tablets are indicated for the palliative treatment of advanced carcinoma of the breast or endometrium (i.e., recurrent, inoperable, or metastatic disease). It should not be used in lieu of currently accepted procedures such as surgery, radiation, or chemotherapy.

CONTRAINDICATIONS

History of hypersensitivity to megestrol acetate or any component of the formulation.

WARNINGS

Megestrol acetate may cause fetal harm when administered to a pregnant woman. Fertility and reproduction studies with high doses of megestrol acetate have shown a reversible feminizing effect on some male rat fetuses. There are no adequate and well-controlled studies in pregnant women. If this drug is used during pregnancy, or if the patient becomes pregnant while taking (receiving) this

drug, the patient should be apprised of the potential hazard to the fetus. Women of childbearing potential should be advised to avoid becoming pregnant.

The use of megestrol in other types of neoplastic disease is not recommended.

(See also Precautions: Carcinogenesis, Mutagenesis, and Impairment of Fertility section.)

The glucocorticoid activity of megestrol acetate tablets has not been fully evaluated. Clinical cases of new onset diabetes mellitus, exacerbation of pre-existing diabetes mellitus, and overt Cushing's syndrome have been reported in association with the chronic use of megestrol. In addition, clinical cases of adrenal insufficiency have been observed in patients receiving or being withdrawn from chronic megestrol therapy in the stressed and non-stressed state. Furthermore, adrenocorticotropin (ACTH) stimulation testing has revealed the frequent occurrence of asymptomatic pituitary-adrenal suppression in patients treated with chronic megestrol therapy. Therefore, the possibility of adrenal insufficiency should be considered in any patient receiving or being withdrawn from chronic megestrol therapy who presents with symptoms and/or signs suggestive of hypoadrenalism (e.g., hypotension, nausea, vomiting, dizziness, or weakness) in either the stressed or non-stressed state. Laboratory evaluation for adrenal insufficiency and consideration of replacement or stress doses of a rapidly acting glucocorticoid are strongly recommended in such patients. Failure to recognize inhibition of the hypothalamic-pituitary-adrenal axis may result in death. Finally, in patients who are receiving or being withdrawn from chronic megestrol therapy, consideration should be given to the use of empiric therapy with stress doses of a rapidly acting glucocorticoid in conditions of stress or serious intercurrent illness (eg., surgery, infection).

PRECAUTIONS

General

Close surveillance is indicated for any patient treated for recurrent or metastatic cancer. Use with caution in patients with a history of thromboembolic disease.

Use in Diabetics

Exacerbation of pre-existing diabetes with increased insulin requirements has been reported in association with the use of megestrol acetate.

Information for the Patients

Patients using megestrol acetate should receive the following instructions:

1. This medication is to be used as directed by the physician.
2. Report any adverse reaction experiences while taking this medication.

Laboratory Tests

Breast malignancies in which estrogen and/or progesterone receptors are positive are more likely to respond to megestrol.

Carcinogenesis, Mutagenesis, and Impairment of Fertility

Administration of megestrol acetate to female dogs for up to 7 years is associated with an increased incidence of both benign and malignant tumors of the breast. Comparable studies in rats and studies in monkeys are not associated with an increased incidence of tumors. The relationship of the dog tumors to humans is unknown but should be considered in assessing the benefit-to-risk ratio when prescribing megestrol acetate and in surveillance of patients on therapy (see Warnings section).

Pregnancy

Pregnancy Category D

See Warnings section.

Nursing Mothers

Because of the potential for adverse effects on the newborn, nursing should be discontinued if megestrol is required for treatment of cancer.

Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

Geriatric Use

Insufficient data from clinical studies of megestrol acetate tablets are available for patients 65 years of age and older to determine whether they respond differently than younger patients. Other reported clinical experience has not identified differences in responses between elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

Megestrol acetate is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

ADVERSE REACTIONS

Weight Gain

Weight gain is a frequent side effect of megestrol. This gain has been associated with increased appetite and is not necessarily associated with fluid retention.

Thromboembolic Phenomena

Thromboembolic phenomena including thrombophlebitis and pulmonary embolism (in some cases fatal) have been reported.

Glucocorticoid Effects

See Warnings section.

Other Adverse Reactions

Heart failure, nausea and vomiting, edema, breakthrough menstrual bleeding, dyspnea, tumor flare (with or without hypercalcemia), hyperglycemia, glucose intolerance, alopecia, hypertension, carpal tunnel syndrome, mood changes, hot flashes, malaise, asthenia, lethargy, sweating and rash.

OVERDOSAGE

No serious unexpected side effects have resulted from studies involving megestrol acetate administered in dosages as high as 1600 mg/day. Oral administration of large, single doses of megestrol acetate (5 g/kg) did not produce toxic effects in mice. Megestrol acetate has not been tested for dialyzability; however, due to its low solubility it is postulated that this would not be an effective means of treating overdose.

DOSAGE AND ADMINISTRATION

Breast cancer: 160 mg/day (40 mg q.i.d.).

Endometrial carcinoma: 40 mg/day to 320 mg/day in divided doses.

At least 2 months of continuous treatment is considered an adequate period for determining the efficacy of megestrol acetate.

HOW SUPPLIED

Megestrol Acetate Tablets, USP

20 mg white, round, scored tablets

(Identified 54 763)

NDC 0054-8603-25: Unit dose, 10 tablets per strip, 10 strips per shelf pack, 10 shelf packs per shipper.

NDC 0054-4603-25: Bottles of 100 tablets.

40 mg white, round, scored tablets

(Identified 54 352)

NDC 0054-8604-25: Unit dose, 10 tablets per strip, 10 strips per shelf pack, 10 shelf packs per shipper.

NDC 0054-4604-25: Bottles of 100 tablets.

Dispense in a well-closed container

as defined in the USP/NF.

STORAGE

Store at 25°C (77°F); excursions permitted to 15° - 30°C (59° - 86°F) [see USP Controlled Room Temperature]. Protect from temperatures above 40°C (104°F).

SPECIAL HANDLING

Health Hazard Data

There is no threshold limit value established by OSHA, NIOSH, or ACGIH.

Exposure or "overdose" at levels approaching recommended dosing levels could result in side effects described above (see Warnings and Adverse Reactions sections). Women at risk of pregnancy should avoid such exposure.

4056065//04

Revised February 2007

© RLI, 2007


PACKAGE LABEL - MEGESTROL ACETATE TABLETS, USP

Rx Only

Roxane Laboratories, Inc.

20 mg

NDC 0054-8603-25: Unit dose.



N
5 0054-8603-25 2

NDC 0054-8603-25 10 x 10 Unit Dose Tablets

**MEGESTROL
ACETATE
Tablets USP**

20 mg

Each tablet contains 20 mg megestrol acetate.
Usual Dosage: See Package Insert For Complete Prescribing Information.
Store at 25°C (77°F); excursions permitted to 15°-30°C (59°-86°F) [see USP Controlled Room Temperature].
Protect from temperatures above 40°C (104°F).
This unit-dose package is not child-resistant.
If dispensed for outpatient use,
a child-resistant container should be utilized.

R_x only

TABLETS IDENTIFIED 54 763


Roxane Laboratories, Inc.
Columbus, Ohio 43216

NDC 0054-8603-25 10 x 10 Unit Dose Tablets

**MEGESTROL
ACETATE
Tablets USP**

20 mg

LOT
EXP.

 Hochsteger Ingelheim
Roxane Laboratories 4267100/04
© RLI, 2007

NDC 0054-4603-25: Bottles of 100 tablets.

Usual Dosage: (See Package Insert for Complete Prescribing Information.)
Store at 25°C (77°F); excursions permitted to 15°-30°C (59°-86°F) [see USP Controlled Room Temperature].
Protect from temperatures above 40°C (104°F).
Dispense in a well-closed container as defined in the USP NF.
TABLETS IDENTIFIED 54 763

Roxane Laboratories, Inc.
Columbus, Ohio 43216

NDC 0054-4603-25 100 Tablets

**MEGESTROL
ACETATE
Tablets USP**

20 mg

Each tablet contains 20 mg megestrol acetate.
R_x only

 Hochsteger Ingelheim
Roxane Laboratories

EXP. LOT

0054-4603-25

4180115/04
© RLI, 2007

40 mg

NDC 0054-8604-25: Unit dose.



HDC 0054-8604-25 10 x 10 Unit Dose Tablets

**MEGESTROL
ACETATE
Tablets USP**

40 mg

Each tablet contains 40 mg megestrol acetate.

Usual Dosage: See Package Insert For
Complete Prescribing Information.

Store at 25°C (77°F); excursions permitted
to 15°-30°C (59°-86°F) [see USP
Controlled Room Temperature].
Protect from temperatures above 40°C (104°F).

This unit-dose package is not child-resistant.
If dispensed for outpatient use,
a child-resistant container should be utilized.

R_x only

TABLETS IDENTIFIED 54 352

Roxane Laboratories, Inc.
Columbus, Ohio 43216

HDC 0054-8604-25 10 x 10 Unit Dose Tablets

**MEGESTROL
ACETATE
Tablets USP**

40 mg

LOT

EXP.



Reethi ger ingelheim
Roxane Laboratories

4267110W03
© RLI, 2007

NDC 0054-4604-25: Bottles of 100 tablets.

Usual Dosage: See Package
Insert for Complete
Prescribing Information.

Store at 25°C (77°F);
excursions permitted to
15°-30°C (59°-86°F) [see
USP Controlled Room
Temperature]. Protect from
temperatures above
40°C (104°F).

Dispense in a well-closed
container as defined in the
USPINF.

TABLETS IDENTIFIED
54 352

Roxane Laboratories, Inc.
Columbus, Ohio 43216

HDC 0054-4604-25 100 Tablets

**MEGESTROL
ACETATE
Tablets USP**

40 mg

Each tablet contains 40 mg
megestrol acetate.

R_x only



Reethi ger ingelheim
Roxane Laboratories



LOT
EXP.

4190120W03
© RLI, 2007